



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# THE ANALYST.

VOL. V.

JULY, 1878.

No. 4.

*EVALUATION OF ELLIPTIC FUNCTIONS OF THE SECOND AND THIRD SPECIES, ANALYST, VOL. V, PAGES 18 & 19.*

BY CHAS. H. KUMMELL, ASSIST. U. S. LAKE SURV. DETROIT, MICH.

4. FOR the second species and ascending scale of moduli we assume the fundamental form :

$$\int_0^\phi d\varphi \sqrt{[a^2 - c^2 \sin^2 \varphi]} = \int_0^\phi a \mathcal{A}(\varphi) d\varphi = a E_0^\phi \left( \frac{c}{a} \right). \quad (21)$$

From (11) follows:

$$\begin{aligned} \tan \varphi &= \frac{a \sin 2\varphi'}{c + a \cos 2\varphi'}; \\ \therefore \sin \varphi &= \frac{a \sin 2\varphi'}{\sqrt{[c^2 + 2ac \cos 2\varphi' + a^2]}} \\ &= \frac{a \sin \varphi' \cos \varphi'}{\sqrt{[a'^2 - c'^2 \sin^2 \varphi']}} \\ &= \frac{a \sin \varphi' \cos \varphi'}{a' \mathcal{A}'(\varphi')}, \end{aligned} \quad (22)$$

$$\begin{aligned} a \mathcal{A}(\varphi) &= \frac{a}{a' \mathcal{A}'(\varphi')} \sqrt{(a'^2 - c'^2 \sin^2 \varphi' - c^2 \sin^2 \varphi' \cos^2 \varphi')} \\ &= \frac{a}{a' \mathcal{A}'(\varphi')} (a' - c \sin^2 \varphi') \\ &= a' \mathcal{A}'(\varphi') + \frac{\frac{1}{2}(a^2 - c^2)}{a' \mathcal{A}'(\varphi')}, \end{aligned} \quad (23)$$

consequently regarding (12)

$$\begin{aligned} a \mathcal{A}(\varphi) d\varphi &= a^2 \mathcal{A}(\varphi)^2 \cdot \frac{d\varphi}{a \mathcal{A}(\varphi)} \\ &= \left[ a' \mathcal{A}'(\varphi') + \frac{\frac{1}{2}(a^2 - c^2)}{a' \mathcal{A}'(\varphi')} \right]^2 \frac{d\varphi'}{a' \mathcal{A}'(\varphi')} \\ &= a' \mathcal{A}'(\varphi') d\varphi' + \frac{1}{2}(a^2 - c^2) \frac{d\varphi'}{a' \mathcal{A}'(\varphi')} + \frac{1}{16}(a^2 - c^2)^2 \frac{d\varphi'}{a'^3 \mathcal{A}'(\varphi')^3} \end{aligned} \quad (24)$$

Differentiating (22) we have

$$\begin{aligned} a\varphi \cos \varphi &= \frac{ad\varphi'}{a'^3 D'(\varphi')^3} \left[ a'^2(1 - 2\sin^2 \varphi') + c'^2 \sin^4 \varphi' \right] \\ &= \frac{ad\varphi'}{a'^3 D'(\varphi')^3} \left[ \frac{a'^4 D'(\varphi')^4}{c'^2} + a'^2 - \frac{a'^4}{c'^2} \right] \\ &= \frac{d\varphi'}{ca'^3 D'(\varphi')^3} \left[ a'^4 D'(\varphi')^4 - \frac{1}{16}(a^2 - c^2)^2 \right], \end{aligned} \quad (25)$$

$$\therefore \frac{1}{16}(a^2 - c^2)^2 \frac{d\varphi'}{a'^3 D'(\varphi')^3} = a' D'(\varphi') d\varphi' - c \cos \varphi d\varphi. \quad (25')$$

Adding this to (24) and integrating we obtain

$$\int_0^\phi a D(\varphi) d\varphi = 2 \int_0^{\phi'} a' D'(\varphi) d\varphi + a'(a - c) \int_0^{\phi'} \frac{d\varphi}{a' D'(\varphi)} - c \sin \varphi$$

$$\text{or} \quad a E_0^\phi \left( \frac{c}{a} \right) = 2a' E_0^{\phi'} \left( \frac{c'}{a'} \right) + a'(a - c) \cdot \frac{1}{a'} F_0^{\phi'} \left( \frac{c'}{a'} \right) - c \sin \varphi. \quad (26)$$

At the next higher step in the scale of moduli we have

$$2a' E_0^{\phi'} \left( \frac{c'}{a'} \right) = 2^2 a'' E_0^{\phi''} \left( \frac{c''}{a''} \right) + 2a''(a' - c') \cdot \frac{1}{a''} F_0^{\phi''} \left( \frac{c''}{a''} \right) - 2c' \sin \varphi'. \quad (26')$$

Finally when

$$a^{(n)} = c^{(n)} = \left\| \frac{1}{2} \left\{ \begin{matrix} a & + & c \\ & \times & \end{matrix} \right\}^{\frac{1}{2}} \right\| \quad \text{we have}$$

$$2^n a^{(n)} E_0^{\phi^{(n)}}(1) = 2^n a^n \sin \varphi^{(n)}. \quad (26'')$$

Adding (26), (26'), ... (26'') we have regarding (13)

$$\begin{aligned} a E_0^\phi \left( \frac{c}{a} \right) &= 2^n a^{(n)} \sin \varphi^{(n)} + \sum_0^{n-1} [2^r a^{(r+1)} (a^{(r)} - c^{(r)})] \cdot \frac{1}{a^{(n)}} \log \tan \frac{1}{2}(\frac{1}{2}\pi + \varphi^{(n)}) \\ &\quad - \sum_0^{n-1} [2^r c^{(r)} \sin \varphi^{(r)}]. \end{aligned} \quad (27)$$

This is the length of an arc of an ellipse the equation of which is

$$\frac{x^2}{a^2} + \frac{y^2}{a^2 - c^2} = 1.$$

between  $x = 0$  and  $x = a \sin \varphi$ .

5. For the second species and descending scale we take the form :

$$\int_0^\phi d\varphi \sqrt{a^2 \cos^2 \varphi + b^2 \sin^2 \varphi} = \int_0^\phi a D(\varphi) d\varphi = a E_0^\phi \sqrt{1 - \frac{b^2}{a^2}}. \quad (28)$$

Consistent with (15) we have also

$$\sin(2\varphi - \varphi_1) = \frac{a-b}{a+b} \sin \varphi_1 = \cos \varphi_1 (\sin 2\varphi - \tan \varphi_1 \cos 2\varphi); \quad (29)$$

for if we add and then subtract the identity

$$\sin \varphi_1 = \sin \varphi_1$$

and divide the difference by the sum (15) results.

Solving for  $\cos 2\varphi$  we have



Formula (34) might have been also derived from (26) by regarding the following relations, which exist between the quantities with primes and those with subscripts if each series is continued in both directions. We have then the amplitudes

$\varphi^{(n)} = \varphi_{-n}; \dots \varphi' = \varphi_{-1}; \varphi = \varphi; \varphi_i = \varphi_1; \varphi_{ii} = \varphi_2; \dots \varphi_{(n)} = \varphi_n; \quad (37)$   
also the moduli

$$\frac{c^{(n)}}{a^{(n)}} = \left\{ 1 - \frac{b_{-n}^2}{a_{-n}^2} \right\}^{\frac{1}{2}}; \dots \frac{c'}{a'} = \left\{ 1 - \frac{b_{-1}^2}{a_{-1}^2} \right\}^{\frac{1}{2}}; \frac{c}{a} = \left\{ 1 - \frac{b^2}{a^2} \right\}^{\frac{1}{2}};$$

$$\frac{c_i}{a_i} = \left\{ 1 - \frac{b_1^2}{a_1^2} \right\}^{\frac{1}{2}}; \dots \frac{c_{(n)}}{a_{(n)}} = \left\{ 1 - \frac{b_n^2}{a_n^2} \right\}^{\frac{1}{2}}; \quad (38)$$

whence

$$\Delta^{(n)}(\varphi) = \Delta_{-n}(\varphi); \dots \Delta'(\varphi) = \Delta_{-1}(\varphi); \Delta\varphi = \Delta\varphi; \Delta_i(\varphi) = \Delta_1(\varphi);$$

$$\dots \Delta_{(n)}(\varphi) = \Delta_n(\varphi). \quad (39)$$

By (12) we have

$$\int_0^{\varphi^{(n)}} \frac{d\varphi}{a^{(n)} \Delta^{(n)}(\varphi)} = \int_0^{\varphi^{(n-1)}} \frac{d\varphi}{a^{(n-1)} \Delta^{(n-1)}(\varphi)} = \dots \int_0^{\varphi'} \frac{d\varphi}{a' \Delta'(\varphi)} = \int_0^{\varphi} \frac{d\varphi}{a \Delta(\varphi)}$$

$$= \int_0^{\psi_i} \frac{d\psi^*}{a_i \Delta_i(\psi)} = \dots \int_0^{\psi_{(n)}} \frac{d\psi}{a_{(n)} \Delta_{(n)}(\psi)} \quad (40)$$

while by (16) we have

$$\int_0^{\psi_{-n}} \frac{d\psi}{2^{-n} a_{-n} \Delta_{-n}(\psi)} = \int_0^{\psi_{-n+1}} \frac{d\psi}{2^{-n+1} a_{-n+1} \Delta_{-n+1}(\psi)} = \dots \int_0^{\psi_{-1}} \frac{d\psi}{2^{-1} a_{-1} \Delta_{-1}(\psi)}$$

$$= \int_0^{\psi} \frac{d\psi}{2a \Delta(\psi)} = \int_0^{\psi_1} \frac{d\psi}{2a_1 \Delta_1(\psi)} = \dots \int_0^{\psi_n} \frac{d\psi}{2^n a_n \Delta_n(\psi)} \quad (41)$$

We have then regarding (37) and (39)

$$a^{(n)} = 2^{-n} a_{-n}; \quad a^{(n-1)} = 2^{-n+1} a_{-n+1}; \dots a' = 2^{-1} a_{-1}; \quad a = a; \quad a_i = 2a_1;$$

$$\dots a_{(n)} = 2^n a_n \quad (42)$$

These relations enable us to derive (34) from (26) and vice versa.

We have by (26) for the next lower step in the modular scale

$$\int_0^{\psi} a_i \Delta_i(\psi) d\psi = 2 \int_0^{\psi} a \Delta(\psi) d\psi + \frac{1}{2} (a_i^2 - c_i^2) \int_0^{\psi} \frac{d\psi}{a \Delta(\psi)} - c_i \sin \psi_i,$$

$$\dots \int_0^{\psi} a \Delta(\psi) d\psi = \frac{1}{2} \int_0^{\psi_i} a_i \Delta_i(\psi) d\psi - \frac{1}{4} (a_i^2 - c_i^2) \int_0^{\psi} \frac{d\psi}{a \Delta(\psi)} + \frac{1}{2} c_i \sin \psi_i.$$

But  $\psi_i = \psi_1$ ;  $a_i = 2a_1$ ;  $\Delta_i(\psi) = \Delta_1(\psi)$ ;  $a_i^2 - c_i^2 = 4b_1^2$ ;  $c_i = 2\sqrt{a_1^2 - b_1^2}$

$$= a - b \text{ and } \int_0^{\psi} \frac{d\psi}{a \Delta(\psi)} = \int_0^{\psi_1} \frac{d\psi}{2a_1 \Delta_1(\psi)} \text{ by (41),}$$

$$\dots \int_0^{\psi} a \Delta(\psi) d\psi = \int_0^{\psi_1} a_1 \Delta_1(\psi) d\psi - \frac{1}{2} b_1^2 \int_0^{\psi_1} \frac{d\psi}{a_1 \Delta_1(\psi)} + \frac{1}{2} (a - b) \sin \psi_1,$$

which is (34).

[\*For want of sorts,  $\psi$  is here, and throughout the rest of this article, used for  $\phi$ —Comp.]

These relations are also very useful for the inverse problem of determining the amplitude to a given integral. It is not my purpose however to show this now.

6. For the third species and ascending scale we take the fundamental form:

$$\int_0^\psi \frac{d\phi}{(p+r\sin^2\phi)\sqrt{[a^2-c^2\sin^2\phi]}} = \int_0^\psi \frac{d\phi}{aP(\phi)\Delta(\phi)} = \frac{1}{ap} \Pi_0^\psi\left(\frac{c}{a}, \frac{r}{p}\right). \quad (43)$$

We have, substituting (22)

$$\frac{1}{P(\phi)} = \frac{a'^2 - c'^2 \sin^2 \phi'}{a'^2 p + a(ar - cp) \sin^2 \phi' - a^2 r \sin^4 \phi'} \quad (44)$$

Assume

$$a'^4 p^2 + aa'^2 p(ar - cp) \sin^2 \phi' - a^2 a'^2 pr \sin^4 \phi' = (a'^2 p + am' \sin^2 \phi')(a'^2 p + am' \sin^2 \phi')$$

$$\text{whence} \quad m' + m' = ar - cp; \quad m'm' = -a'^2 pr \quad (45)$$

$$\text{and} \quad m' - m' = \sqrt{(p+r)(ar^2 + cp^2)}. \quad (46)$$

Now let

$$\begin{aligned} \frac{\frac{1}{p} - \frac{ac}{a'^2 p} \sin^2 \phi'}{\left(1 + \frac{am'}{a'^2 p} \sin^2 \phi'\right) \left(1 + \frac{am'}{a'^2 p} \sin^2 \phi'\right)} &= \frac{1}{p' + r' \sin^2 \phi'} + \frac{1}{p' + r' \sin^2 \phi'} \\ &= \frac{1}{p'} + \frac{1}{p'} + \frac{r' + r'}{p' p'} \sin^2 \phi' \div \left(1 + \frac{r'}{p'} \sin^2 \phi'\right) \left(1 + \frac{r'}{p'} \sin^2 \phi'\right). \end{aligned}$$

By comparison and regarding (45) we have

$$r' = \frac{am' p'}{a'^2 p} = -\frac{p'}{m'} ar; \quad r' = \frac{am' p'}{a'^2 p} = -\frac{p'}{m'} ar. \quad (47)$$

$$\frac{1}{p} = \frac{1}{p'} + \frac{1}{p'}; \quad -c = \frac{m'}{p'} + \frac{m'}{p'} \quad (48)$$

$$\text{whence} \quad p' = \frac{m' - m'}{m' + cp} p; \quad p' = -\frac{m' - m'}{m' + cp} p. \quad (49)$$

Thus  $p'$ ,  $p'$  and  $r'$ ,  $r'$  are known, and placing  $P'(\phi) = p' + r' \sin^2 \phi$ ;  $P(\phi) = p' + r' \sin^2 \phi$  we have regarding (12) or (41)

$$\int_0^\psi \frac{d\phi}{aP(\phi)\Delta(\phi)} = \int_0^\psi \frac{d\phi}{a'P'(\phi)\Delta'(\phi)} + \int_0^\psi \frac{d\phi}{a'P''(\phi)\Delta'(\phi)} \quad (50)$$

At the next higher step in the modular scale we have besides (11') the formulæ

$$m'' + m'' = a'r' - c'p', \quad m'' + m'' = a'r' - c'p', \quad (45')$$

$$m'' - m'' = \sqrt{[(p' + r')(a'^2 r' + c'^2 p')]}, \quad m'' - m'' = \sqrt{[(p' + r')(a'^2 r' + c'^2 p')]}, \quad (46')$$

$$p'' = \frac{m'' - m''}{m'' + c'p'} p'; \quad p'' = -\frac{m'' - m''}{m'' + c'p'} p' \quad p'' = \frac{m'' - m''}{m'' + c'p'} p'; \quad p'' = -\frac{m'' - m''}{m'' + c'p'} p' \quad \dots (49')$$

$$r'' = -\frac{p''}{m'} a' r'; \quad r' = -\frac{p'}{m''} a' r'; \quad r'' = -\frac{p''}{m''} a' r'; \quad r' = -\frac{p'}{m'} a' r'; \quad (47')$$

and if  $P''(\phi) = p'' + r'' \sin^2 \phi$ ;  $P'(\phi) = p' + r' \sin^2 \phi$ ;  $P(\phi) = p + r \sin^2 \phi$ ;  $P''(\phi) = p'' + r'' \sin^2 \phi$  we have

$$\int_0^\psi \frac{d\phi}{a P(\phi) \Delta(\phi)} = \int_0^{\phi'} \frac{d\phi}{a' P'(\phi) \Delta'(\phi)} = \int_0^{\phi''} \frac{d\phi}{a'' P''(\phi) \Delta''(\phi)} + \int_0^{\phi'} \frac{d\phi}{a'' P''(\phi) \Delta''(\phi)} + \int_0^{\phi''} \frac{d\phi}{a'' P''(\phi) \Delta''(\phi)} + \int_0^{\phi''} \frac{d\phi}{a'' P''(\phi) \Delta''(\phi)} \quad (51)$$

Pursuing the ascending scale of moduli until  $a^{(n)} = c^{(n)} = \left\| \frac{1}{2} \left( a + c \right) \right\|$  and consequently  $\Delta^{(n)}(\phi) = \cos \phi$ , we have finally  $2^n$  integrals of the form:

$$\begin{aligned} \int_0^{\phi^{(n)}} \frac{d\phi}{a^{(n)} P^{(n)}(\phi) \cos \phi} &= \frac{r^{(n)}}{a^{(n)}(p^{(n)} + r^{(n)})} \int_0^{\phi^{(n)}} \frac{d\phi \cos \phi}{P^{(n)}(\phi)} + \frac{1}{a^{(n)}(p^{(n)} + r^{(n)})} \int_0^{\phi^{(n)}} \frac{d\phi}{\cos \phi} \\ &= \frac{1/r^{(n)}}{a^{(n)}(p^{(n)} + r^{(n)}) \sqrt{p^{(n)}}} \tan^{-1} \left\{ \sqrt{\left( \frac{r^{(n)}}{p^{(n)}} \right) \sin \phi^{(n)}} \right\} \\ &\quad + \frac{1}{a^{(n)}(p^{(n)} + r^{(n)})} \log \tan \frac{1}{2} \left( \frac{1}{2} \pi + \phi^{(n)} \right). \end{aligned} \quad (51)$$

These reductions are very troublesome if  $p', p'', \dots, r', r'', \dots$  are complex quantities. In this case and if a problem leads to the form

$$\int_0^\phi f(\sin^2 \phi) \frac{d\phi}{a \Delta(\phi)}$$

( $f$  denoting a rational function of  $\sin^2 \phi$ ) which might be expressed by means of some of the three species of elliptic functions and elementary forms, it is perhaps better to express  $f(\sin^2 \phi)$  successively in terms of  $\sin^2 \phi', \sin^2 \phi'', \dots, \sin^2 \phi^{(n)}$  so that if  $f'(\sin^2 \phi), f''(\sin^2 \phi), \dots$  are the resulting forms we have

$$\begin{aligned} \int_0^\phi f(\sin^2 \phi) \frac{d\phi}{a \Delta(\phi)} &= \int_0^{\phi'} f'(\sin^2 \phi) \frac{d\phi}{a' \Delta'(\phi)} = \int_0^{\phi''} f''(\sin^2 \phi) \frac{d\phi}{a'' \Delta''(\phi)} \dots \\ &= \int_0^{\phi^{(n)}} f^{(n)}(\sin^2 \phi) \frac{d\phi}{a^{(n)} \cos \phi} = \frac{1}{a^{(n)}} \int_0^{\sin \phi^{(n)}} \frac{f(x^2) dx}{1 - x^2}. \end{aligned} \quad (52)$$

7. For the third species and descending scale we take the form:

$$\begin{aligned} \int_0^\phi \frac{d\phi}{p \cos^2 \phi + q \sin^2 \phi} \sqrt{[a^2 \cos^2 \phi + b^2 \sin^2 \phi]} &= \int_0^\phi \frac{d\phi}{a P(\phi) \Delta(\phi)} \\ &= \Pi_0^\phi \left\{ \sqrt{\left( 1 - \frac{b^2}{a^2} \right)}, \frac{p}{q} - 1 \right\}. \end{aligned} \quad (53)$$

Substituting (31) we have regarding (16) or (41)

$$\frac{d\phi}{a P(\phi) \Delta(\phi)} = \frac{1}{p + \frac{1}{2}(q - p) \left[ 1 + \frac{a - b}{a + b} \sin^2 \psi_1 - \cos \varphi_1 \Delta_1(\psi_1) \right]} \frac{d\phi_1}{2a_1 \Delta_1(\psi)}$$

$$\begin{aligned}
 &= \frac{p + \frac{1}{2}(q-p) \left[ 1 + \frac{a-b}{a+b} \sin^2 \psi_1 + \cos^2 \psi_1 \Delta_1(\psi) \right]}{pq + \frac{q-p}{(a+b)^2} (a^2 q - b^2 p) \sin^2 \psi_1} \cdot \frac{d\psi_1}{2a_1 \Delta_1(\psi_1)} \\
 &= \frac{a^2 q^2 - b^2 p^2}{4(a^2 q - b^2 p)} \cdot \frac{1}{pq \cos^2 \psi_1 + \left( \frac{aq+bp}{a+b} \right)^2 \sin^2 \psi_1} \cdot \frac{d\psi_1}{a_1 \Delta_1(\psi_1)} \\
 &+ \frac{a^2 - b^2}{4(a^2 q - b^2 p)} \cdot \frac{d\psi_1}{a_1 \Delta_1(\psi_1)} + \frac{q-p}{4a_1} \cdot \frac{d\psi_1 \cos \psi_1}{pq \cos^2 \psi_1 + \left( \frac{aq+bp}{a+b} \right)^2 \sin^2 \psi_1}. \quad (54)
 \end{aligned}$$

Let 
$$m_1 = \frac{a^2 q^2 - b^2 p^2}{a^2 q - b^2 p}, \quad (55)$$

$$p_1 = \frac{q}{m_1} p, \quad (56)$$

$$q_1 = \frac{(aq+bp)^2}{4a_1^2 m_1}. \quad (57)$$

We have then if  $P_1(\psi_1) = p_1 \cos^2 \psi_1 + q_1 \sin^2 \psi_1$

$$\begin{aligned}
 \int_0^\psi \frac{d\psi}{aP(\psi)\Delta(\psi)} &= \frac{1}{4} \left\{ \int_0^{\psi_1} \frac{d\psi}{a_1 P_1(\psi)\Delta_1(\psi)} + \frac{a^2 - b^2}{a^2 q - b^2 p} \int_0^{\psi_1} \frac{d\psi}{a_1 \Delta_1(\psi)} \right. \\
 &\quad \left. + \frac{q-p}{a_1 m_1} \int_0^{\psi_1} \frac{d\psi \cos \psi}{P_1(\psi)} \right\};
 \end{aligned}$$

or if we put  $q-p = r; q_1 - p_1 = r_1$  (58)

we have  $r_1 = \frac{a^2 q - b^2 p}{4a_1^2 m_1} r$ ; hence  $\frac{a^2 - b^2}{a^2 q - b^2 p} = \frac{(a-b)r}{2a_1 m_1 r_1}$  and

$$\begin{aligned}
 \int_0^\psi \frac{d\psi}{aP(\psi)\Delta(\psi)} &= \frac{1}{4} \left\{ \int_0^{\psi_1} \frac{d\psi}{a_1 P_1(\psi)\Delta_1(\psi)} + \frac{(a-b)r}{2a_1 m_1 r_1} \int_0^{\psi_1} \frac{d\psi}{a_1 \Delta_1(\psi)} \right. \\
 &\quad \left. + \frac{r}{a_1 m_1} \int_0^{\psi_1} \frac{d\psi \cos \psi}{P_1(\psi)} \right\}. \quad (59)
 \end{aligned}$$

For the next lower step in the modular scale we have

$$\begin{aligned}
 \frac{1}{4} \int_0^{\psi_1} \frac{d\psi}{a_1 P_1(\psi)\Delta_1(\psi)} &= \frac{1}{4^2} \left\{ \int_0^{\psi_2} \frac{d\psi}{a_2 P_2(\psi)\Delta_2(\psi)} + \frac{(a_1 - b_1)r_1}{2a_2 m_2 r_2} \int_0^{\psi_2} \frac{d\psi}{a_2 \Delta_2(\psi)} \right. \\
 &\quad \left. + \frac{r_1}{a_2 m_2} \int_0^{\psi_2} \frac{d\psi \cos \psi}{P_2(\psi)} \right\}, \quad (59_1)
 \end{aligned}$$

where we have besides (17)

$$m_2 = \frac{a_1^2 q_1^2 - b_1^2 p_1^2}{a_1^2 q_1 - b_1^2 p_1}, \quad (55_1)$$

$$p_2 = \frac{q_1}{m_2} p_1, \quad (56_1)$$



$$q_2 = \frac{(a_1 q_1 + b_1 p_1)^2}{4a_2^2 m_2}, \quad (57_1)$$

$$r_2 = \frac{a_1^2 q_1 - b_1^2 p_1}{4a_2^2 m_2} r_1 = q_2 - p_2. \quad (58_1)$$

Descending the modular scale until  $a_n = b_n = \left\| \frac{1}{2}(a \times b) \right\|$  we have finally

$$\frac{1}{4^n} \int_0^{\psi_n} \frac{d\phi}{a_n P_n(\phi)} = \frac{1}{4^n a_n \sqrt{(p_n q_n)}} \tan^{-1} \left\{ \sqrt{\left( \frac{q_n}{p_n} \right)} \tan \phi_n \right\}. \quad (59_n)$$

Adding (59), (59<sub>1</sub>), ... (59<sub>n</sub>) we obtain

$$\begin{aligned} \int_0^{\psi} \frac{d\phi}{a P(\phi) \Delta(\phi)} &= \frac{1}{4^n a_n \sqrt{(p_n q_n)}} \tan^{-1} \left\{ \left( \frac{q_n}{p_n} \right)^{\frac{1}{2}} \tan \phi_n \right\} + \sum_1^n \left\{ \frac{(a_{s-1} - b_{s-1}) r_{s-1}}{2^{2s+1} a_s m_s r_s} \right. \\ &\quad \times \left. \int_0^{\psi_s} \frac{d\phi}{a_s \Delta_s(\phi)} \right\} + \sum_1^n \left\{ \frac{r_{s-1}}{2^{2s} a_s m_s} \int_0^{\psi_s} \frac{d\phi \cos \phi}{P_s(\phi)} \right\}. \quad (60) \end{aligned}$$

$$\text{But} \quad \int_0^{\phi_s} \frac{d\phi}{a_s \Delta_s(\phi)} = \frac{\phi_s}{2^{n-s} a_n} \quad (61)$$

$$\text{and} \quad \int_0^{\phi_s} \frac{d\phi \cos \phi}{P_s(\phi)} = \frac{1}{\sqrt{(p_s r_s)}} \tan^{-1} \left\{ \left( \frac{r_s}{p_s} \right)^{\frac{1}{2}} \sin \phi_s \right\}, \quad (62)$$

consequently

$$\begin{aligned} \int_0^{\psi} \frac{d\phi}{a P(\phi) \Delta(\phi)} &= \frac{\tan^{-1} \left[ \sqrt{\left( \frac{q_n}{p_n} \right)} \tan \phi_n \right]}{2^{2n} a_n \sqrt{(p_n q_n)}} + \sum_1^n \left\{ \frac{(a_{s-1} - b_{s-1}) r_{s-1}}{2^{n+s+1} a_s m_s r_s} \right\} \frac{\phi_n}{a_n} \\ &\quad + \sum_1^n \left\{ \frac{r_{s-1}}{2^{2s} a_s m_s \sqrt{(p_s r_s)}} \tan^{-1} \left[ \left( \frac{r_s}{p_s} \right)^{\frac{1}{2}} \sin \phi_s \right] \right\}. \quad (63) \end{aligned}$$

It is hardly necessary to state that if  $p_s$  has the opposite sign to  $q_s$  or  $r_s$ , the  $\tan^{-1}$  must be replaced by logarithms.

If  $\phi = \frac{1}{2}\pi$  then  $\phi_1 = \pi$ ;  $\phi_2 = 2\pi$ ; ...  $\phi_n = 2^{n-1}\pi$ ; hence

$$\int_0^{\frac{\pi}{2}} \frac{d\phi}{a P(\phi) \Delta(\phi)} = \frac{\pi}{2^{n+1} a_n \sqrt{(p_n q_n)}} + \sum_1^n \left\{ \frac{(a_{s-1} - b_{s-1}) r_{s-1}}{2^{s+2} a_s m_s r_s} \right\} \frac{\pi}{a_n}, \quad (64)$$

for

$$\int_0^{2^{s-1}\pi} \frac{d\phi \cos \phi}{P_s(\phi)} = 0 = \int_0^{\pi} \frac{d\phi \cos \phi}{P_s(\phi)},$$

because the elements of this integral from 0 to  $\frac{1}{2}\pi$  are equal and opposite in sign to those from  $\frac{1}{2}\pi$  to  $\pi$ .

Equation (59) could be inverted by means of the relations (38), ... (42). I have not yet succeeded however in reducing the relation between the  $p$ ,  $q$  and  $r$  to a convenient form. The reduction by ascending the modular scale as made under No. 6 seems to be preferable.